

Trigonometry - Angles and Ratios Lesson #2: Trigonometric Ratios for Angles from 0° to 360°


Pythagorean Theorem

The traditional formula for the Pythagorean Theorem is $c^2 = a^2 + b^2$.

In trigonometry, we use x , y , and r instead of a , b , and c .

The point $P(x, y)$ lies on the terminal arm of angle θ .

The distance from the origin to point P is r , the radius of the circle formed by the rotation.



$$x^2 + y^2 = r^2, \text{ where } r > 0$$


Standard position.

Class Ex. #1



Sketch the rotation angle in standard position, and calculate the exact distance from the origin to the given point. Where appropriate, write the answer in simplest mixed radical form.

- a) Point $P(-5, 12)$ on the terminal arm of angle θ .




$$r^2 = x^2 + y^2$$

$$r = \sqrt{(-5)^2 + (12)^2}$$

$$r = \sqrt{169}$$

$$r = 13$$

- b) Point $Q(-2, -6)$ on the terminal arm of angle A .



$$r^2 = x^2 + y^2$$

$$r = \sqrt{(-2)^2 + (-6)^2}$$

$$r = \sqrt{40}$$

$$r = 2\sqrt{10}$$

Given $\theta = 224^\circ$:

- state the quadrant
- draw the angle in standard position
- state the reference angle
- give one positive & one negative coterminal angle.

Quad III

$$\theta + 360^\circ n$$


$$224^\circ + 360^\circ = 584^\circ$$

$$224^\circ - 360^\circ = -136^\circ$$

ref L

$$= 224^\circ - 180^\circ$$


$$= 44^\circ$$



Trigonometric Ratios

Complete the following:


sine ratio $\Rightarrow \sin \theta = \frac{O}{H}$
cosine ratio $\Rightarrow \cos \theta = \frac{A}{H}$
tangent ratio $\Rightarrow \tan \theta = \frac{O}{A}$



These ratios are called the **Primary Trigonometric Ratios** and can be remembered by the acronym **SOHCAHTOA**.



Write the primary trigonometric ratios for angle θ in terms of x , y , and r .



$\sin \theta = \frac{O}{H} \rightarrow \frac{y}{r}$
 $\cos \theta = \frac{A}{H} \rightarrow \frac{x}{r}$
 $\tan \theta = \frac{O}{A} \rightarrow \frac{y}{x}$




You should memorize these formulas.

Some students use a phrase like “seven yellow rabbits” to remember $\sin \theta = \frac{y}{r}$.



The point $(15, 8)$ lies on the terminal arm of an angle θ as shown. Calculate the value of r , and hence determine the exact values of the primary trigonometric ratios.




x, y, r **3 amigos**
 $r = \sqrt{15^2 + 8^2}$
 $r = \sqrt{289}$
 $r = 17$

Complete Assignment Questions #1 - #5

$\sin \theta = \frac{8}{17}$ $\cos \theta = \frac{15}{17}$ $\tan \theta = \frac{8}{15}$

Part 4

The rotation angle in Part 1 is reflected in the x-axis. Complete the following:



- a) The point $S(x, y)$ has coordinates $S(\quad , \quad)$.
- b) The reference angle is \quad and the rotation angle is \quad .
- c) $\sin 300^\circ = \frac{y}{r} = \quad$ $\cos 300^\circ = \frac{x}{r} = \quad$ $\tan 300^\circ = \frac{y}{x} = \quad$
- d) Confirm these trigonometric ratios on your calculator.

Observations

- The trigonometric ratios for angles between 90° and 360° are either the trigonometric ratios of the reference angle, or the negative of the trigonometric ratios of the reference angle.
- The sign of the trigonometric ratios depends on the quadrant and whether x and y are positive or negative.

Determining the Sign of a Trigonometric Ratio

- a) In quadrant 1, draw the rotation angle θ in standard position and complete the table.
- b) Repeat for quadrants 2 - 4.

Sin	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 2</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{-}$</td><td>$-$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{+}{-}$</td><td>$-$</td></tr> </table>	Quadrant 2			$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{+}{-}$	$-$	$\tan \theta = \frac{y}{x}$	$\frac{+}{-}$	$-$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 1</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> </table>	Quadrant 1			$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{+}{+}$	$+$	All
	Quadrant 2																										
$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{-}$	$-$																									
$\tan \theta = \frac{y}{x}$	$\frac{+}{-}$	$-$																									
Quadrant 1																											
$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{+}{+}$	$+$																									
<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 3</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> </table>	Quadrant 3			$\sin \theta = \frac{y}{r}$	$\frac{-}{-}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{-}{-}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{-}{-}$	$+$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 4</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{-}{+}$</td><td>$-$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{-}{+}$</td><td>$-$</td></tr> </table>	Quadrant 4			$\sin \theta = \frac{y}{r}$	$\frac{-}{+}$	$-$	$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{-}{+}$	$-$		
Quadrant 3																											
$\sin \theta = \frac{y}{r}$	$\frac{-}{-}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{-}{-}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{-}{-}$	$+$																									
Quadrant 4																											
$\sin \theta = \frac{y}{r}$	$\frac{-}{+}$	$-$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{-}{+}$	$-$																									
tan	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 2</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{-}$</td><td>$-$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{+}{-}$</td><td>$-$</td></tr> </table>	Quadrant 2			$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{+}{-}$	$-$	$\tan \theta = \frac{y}{x}$	$\frac{+}{-}$	$-$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 1</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> </table>	Quadrant 1			$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{+}{+}$	$+$	Cos
	Quadrant 2																										
$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{-}$	$-$																									
$\tan \theta = \frac{y}{x}$	$\frac{+}{-}$	$-$																									
Quadrant 1																											
$\sin \theta = \frac{y}{r}$	$\frac{+}{+}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{+}{+}$	$+$																									
<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 3</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{-}{-}$</td><td>$+$</td></tr> </table>	Quadrant 3			$\sin \theta = \frac{y}{r}$	$\frac{-}{-}$	$+$	$\cos \theta = \frac{x}{r}$	$\frac{-}{-}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{-}{-}$	$+$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td colspan="3">Quadrant 4</td></tr> <tr><td>$\sin \theta = \frac{y}{r}$</td><td>$\frac{-}{+}$</td><td>$-$</td></tr> <tr><td>$\cos \theta = \frac{x}{r}$</td><td>$\frac{+}{+}$</td><td>$+$</td></tr> <tr><td>$\tan \theta = \frac{y}{x}$</td><td>$\frac{-}{+}$</td><td>$-$</td></tr> </table>	Quadrant 4			$\sin \theta = \frac{y}{r}$	$\frac{-}{+}$	$-$	$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$	$\tan \theta = \frac{y}{x}$	$\frac{-}{+}$	$-$		
Quadrant 3																											
$\sin \theta = \frac{y}{r}$	$\frac{-}{-}$	$+$																									
$\cos \theta = \frac{x}{r}$	$\frac{-}{-}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{-}{-}$	$+$																									
Quadrant 4																											
$\sin \theta = \frac{y}{r}$	$\frac{-}{+}$	$-$																									
$\cos \theta = \frac{x}{r}$	$\frac{+}{+}$	$+$																									
$\tan \theta = \frac{y}{x}$	$\frac{-}{+}$	$-$																									

c) Complete the following statements using the results from a) and b).

- i) Sine ratios have **positive** values in quadrants 1 and 2.
- ii) Cosine ratios have **positive** values in quadrants 1 and 4.
- iii) Tangent ratios have **positive** values in quadrants 1 and 3.
- iv) Sine ratios have **negative** values in quadrants 3 and 4.
- v) Cosine ratios have **negative** values in quadrants 2 and 3.
- vi) Tangent ratios have **negative** values in quadrants 2 and 4.


CAST Rule

The results can be memorized by:

- the CAST rule or
- by remembering to “Add Sugar To Coffee”

not necessary !!

sin → y
cos → x



Determine, without using technology, whether the given trigonometric ratios are positive or negative.

a) $\sin 340^\circ$ IV
-

b) $\tan 227^\circ$ III
+

c) $\sin 88^\circ$ I
+

d) $\cos 235^\circ$ III
-

e) $\cos 308^\circ$ IV
+

f) $\tan 123^\circ$ II
-

Trigonometric Ratios of an Angle in Terms of the Reference Angle

The trigonometric ratios for any angle are either the trigonometric ratios of the reference angle, or the negative of the trigonometric ratios of the reference angle.

Use the following procedure:

- i) Determine the sign of the ratio (positive or negative).
- ii) Determine the measure of the reference angle.
- iii) Combine i) and ii).

To write $\cos 260^\circ$ as the cosine of an acute angle using the above procedure, we have

- i) negative ii) 80° iii) $\cos 260^\circ = -\cos 80^\circ$.

The result can be verified on a calculator.



Rewrite as the same trigonometric function of an acute angle.

using ref L

a) $\sin 140^\circ$ ⁺
ref L = 40° ⁺ II

b) $\tan 323^\circ$ ⁻
ref L = 37° ⁺ IV

c) $\cos 165^\circ$ ⁻
ref L = 15° ⁺ II

$\sin 140^\circ = \sin 40^\circ$

$\tan 323^\circ = -\tan 37^\circ$

$\cos 165^\circ = -\cos 15^\circ$

d) $\sin 287^\circ$ ⁻
ref L = 73° ⁺ IV

e) $\cos 308^\circ$ ⁺
ref L = 52° ⁺ IV

f) $\tan 199^\circ$ ⁺
ref L = 19° ⁺ III

$\sin 287^\circ = -\sin 73^\circ$

$\cos 308^\circ = \cos 52^\circ$

$\tan 199^\circ = \tan 19^\circ$

Patterns in Trigonometric Ratios

We have the following pattern of results relating the trigonometric ratios of rotation angles to the trigonometric ratios of reference angles.

Let x° be the reference angle for an angle in standard position.

$\sin (180 - x)^\circ = \sin x^\circ$

$\cos (180 - x)^\circ = -\cos x^\circ$

$\tan (180 - x)^\circ = -\tan x^\circ$

$\sin (180 + x)^\circ = -\sin x^\circ$

$\cos (180 + x)^\circ = -\cos x^\circ$

$\tan (180 + x)^\circ = \tan x^\circ$

$\sin (360 - x)^\circ = -\sin x^\circ$

$\cos (360 - x)^\circ = \cos x^\circ$

$\tan (360 - x)^\circ = -\tan x^\circ$

Complete Assignment Questions #6 - #11 and the Group Investigation.

Assignment

#1-13

1. Sketch the rotation angle in standard position, and calculate the exact distance from the origin to the given point.

a) Point $P(15, -8)$ on the terminal arm of angle θ .

b) Point $Q(-24, -7)$ on the terminal arm of angle B .